

Description

Mazuri® Flamingo Breeder can be fed as the staple diet. As the primary portion of the birds' daily food intake, it will maintain the birds' vibrant plumage color when fed according to feeding directions. The diet is applicable to Cuban flamingos, Rody flamingos, cranes and certain other crustacean eating birds.

Features and Benefits

- **Mazuri® Flamingo Breeder floats** - Prevents accumulation on the bottom of the feeding area.
- **Nutritionally balanced** - Contains high levels of vitamins and fish proteins.
- **Contains canthaxanthin** - A pigment within the feed that maintains plumage color.

Product Form

- Extruded pellet size: 5/16" diameter x 3/8" length.

Catalog #0005482

50 lb. net weight paper sack.

Guaranteed Analysis

Crude protein not less than.....	34.0%
Crude fat not less than.....	5.0%
Crude fiber not more than.....	7.0%
Calcium (Ca) not less than	2.60%
Calcium (Ca) not more than.....	3.10%
Phosphorus (P) not less than	1.00%
Sodium (Na) not less than	0.50%
Sodium (Na) not more than	1.00%

Ingredients

Dehydrated alfalfa meal, ground wheat, corn gluten meal, fish meal, brewers dried yeast, porcine meat meal, dehulled soybean meal, wheat flour, calcium carbonate, porcine animal fat preserved with BHA, dicalcium phosphate, monocalcium phosphate, sodium sesquicarbonate, soybean oil, L-lysine, monosodium glutamate, potassium chloride, salt, DL-methionine, menadione dimethylpyrimidinol bisulfite (source of vitamin K), pyridoxine hydrochloride, d-alpha tocopheryl acetate (natural source vitamin E), choline chloride, magnesium oxide, cholecalciferol (vitamin D₃), folic acid, canthaxanthin, vitamin A acetate, L-ascorbyl-2-polyphosphate (vitamin C), biotin, thiamin mononitrate, ethoxyquin (a preservative), calcium pantothenate, riboflavin, vitamin B₁₂ supplement, manganous oxide, inositol, p-aminobenzoic acid, nicotinic acid, calcium iodate, zinc oxide, ferrous carbonate, copper sulfate, zinc sulfate, cobalt carbonate, sodium selenite.

Feeding Directions

Mazuri® Flamingo Breeder can be fed as a major part of the birds' daily food intake during the breeding season, which is during the spring and summer months. At this time the feed should be fed to growing birds and to mature male birds to keep them in peak condition. The feed can be scattered directly on the water, where it will float, or it can be fed in a long, shallow trough, which will prevent the food from floating away from the birds. Adult birds will eat 150-200 grams/day.

Approximate Nutrient Composition**NUTRIENTS**

Protein, %	35.5
Arginine, %.....	1.68
Cystine, %.....	0.45
Glycine, %.....	2.00
Histidine, %.....	0.75
Isoleucine, %.....	1.50
Leucine, %.....	3.20
Lysine, %.....	2.00
Methionine, %.....	1.10
Phenylalanine, %.....	1.60
Tyrosine, %.....	1.21
Threonine, %.....	1.25
Tryptophan, %.....	0.30
Valine, %.....	1.60
Fat, %	5.6
Linoleic Acid, %.....	1.5
Omega-3 Fatty Acids, %.....	0.39
Omega-6 Fatty Acids, %.....	1.48
Fiber (Crude), %	7.0
Neutral Detergent Fiber, %.....	15.0
Acid Detergent Fiber, %.....	9.3
Starch, %.....	16.0
Metabolizable Energy* , kcal/kg.....	2600

MINERALS

Ash, %	11.0
Calcium, %.....	2.78
Phosphorus, %.....	1.29
Phosphorus (non-phytate), %.....	1.06
Potassium, %.....	1.30
Magnesium, %.....	0.23
Sodium, %.....	0.58
Chlorine, %.....	0.68
Sulfur, %.....	0.44
Iron, ppm.....	464
Zinc, ppm.....	105
Manganese, ppm.....	157
Copper, ppm.....	18
Iodine, ppm.....	6.7
Selenium, ppm.....	1.0
Cobalt, ppm.....	1.1

VITAMINS

Thiamin, ppm.....	26
Riboflavin, ppm.....	21
Niacin, ppm.....	51
Pantothenic Acid, ppm.....	37
Choline, ppm.....	1,470
Folic Acid, ppm.....	11
Pyridoxine, ppm.....	21
Biotin, ppm.....	0.40
Vitamin B ₁₂ , mcg/kg.....	96
Vitamin C, ppm.....	90
Vitamin A, IU/kg.....	24,200
Vitamin D ₃ (added), IU/kg.....	4,550
Vitamin E, IU/kg.....	320
Vitamin K (as menadione), ppm.....	7.7
Carotene, ppm.....	12
Canthaxanthin, ppm.....	40

* Calculated using data from domesticated poultry.

Quality Controlled by PMI Nutrition International, a subsidiary of America's oldest and largest animal nutrition company.

Nutrient composition is based on the latest ingredient analysis information. Since nutrient composition of natural ingredients varies, analyses will vary accordingly.